

Gastroenterology Consultants of Long Island

Ian M. Storch, D.O.,FACP,FACG

2001 Marcus Ave. Ste E240
New Hyde Park, NY 11042

Phone (516) 673-4801
Fax: (516) 352-3680

COLONOSCOPY—MOVIPREP

DATE: _____

TIME: _____

****Fill the prescription for the MOVIPREP PREP.**

DAY BEFORE THE COLONOSCOPY

****Stay on a clear liquid diet the ENTIRE DAY (breakfast, lunch, dinner). DO NOT EAT ANY SOLID FOODS OR USE MILK OR MILK PRODUCTS. Acceptable liquids are clear juices, coffee or tea with sugar, clear broths, jello (not red), Italian ices, ice pops, sports drinks, sodas.**

****If preferred, mix solution ahead of time and refrigerate prior to drinking.**

At 5PM empty 1 Pouch A and 1 Pouch B into the disposable Moviprep container. Add lukewarm drinking water to the top line of the container, mix to dissolve powder. Drink 8 oz of Moviprep every 15 minutes until the container is completed.

At 9:00pm empty 1 Pouch A and 1 Pouch B into the disposable Moviprep container. Add lukewarm water to the top line of the container, mix to dissolve powder. Drink 8oz of Moviprep every 15 minutes until the container is completed.

After completing Moviprep fill the empty container up with drinking water. Complete drinking the additional 1 liter of water before midnight to re-hydrate.

It is important that you drink all of the solution and water as instructed. Failure to follow these instructions may negatively affect your procedure.

****CLEAR LIQUIDS CAN BE CONTINUED UNTIL MIDNIGHT—NOTHING TO DRINK AFTER MIDNIGHT OR ON THE MORNING OF THE PROCEDURE**
IF YOU TAKE A HEART MEDICATION OR BLOOD PRESSURE PILL TAKE IT WITH A SMALL AMT. OF WATER ON THE MORNING OF THE TEST.**

*****You are NOT permitted to drive yourself home after your procedure! You must have a responsible adult with you when you arrive for your procedure. You may use a taxi service, car service, or public transportation ONLY if you have a responsible adult with you. Failure to follow this policy will result in cancellation of your procedure EVEN if you have completed a preparation for your test.**

MEDICATIONS TO HOLD FOR THIS PROCEDURE: